

WELCOME TO OUR STD CLINIC



Our goal during clinic is to offer testing and treatment for sexually transmitted infections. Please place a check mark by the tests or treatment you would like today (you can find information about the infections on the back of this page). If you have access the internet, more information on STDs can be found at <http://www.ashastd.org>.

We offer testing for:

- Chlamydia,
- Gonorrhea,
- Syphilis (blood test), and
- HIV (the virus that causes AIDS)*.

We offer treatment for the following STDs:

- Chlamydia,
- Gonorrhea,
- Pelvic Inflammatory Disease,
- Syphilis,
- Vaginal infections (Yeast, Bacterial Vaginosis, Trichomoniasis),
- Genital Warts (HPV),
- Pubic lice (crabs),
- Molluscum Contagiosum, and Scabies.

*Our **HIV testing** can be done with an oral test and your results will be ready today in 20 minutes. We also offer a blood test, but your results will be ready in two weeks and you need to come in person to get your result (no exceptions).



SOME TIPS TO ENSURE THAT YOUR VISIT GOES SMOOTHLY



- **Let the intake person know if you only want an HIV test. This will speed things up.**
- **Be aware that we do not offer Pap smears or screening for Herpes.**
- **Regular testing and treatment are done for free.**
- **We will only contact you if you have an infection that needs treatment.**
- **Please do not use the bathroom until after you see the doctor! If you use the bathroom, we may not be able to do your exam.**

DISEASE	SYMPTOMS	HOW IS IT SPREAD?	TREATMENT
Chlamydia	7-21 days after infection Discharge from penis or vagina Pain or burning with urination Pain in lower abdominal area Maybe NO SYMPTOMS Can lead to Pelvic Inflammatory Disease (PID) if left untreated	Sexual contact (vaginal, oral, anal) with someone who has Chlamydia Mother can pass it to an infant during childbirth	Antibiotics prescribed by a doctor Sexual partners must also be treated
Pubic Lice (crabs)	Immediately to a few days Intense itching in genital area Visual evidence of crabs	Close personal contact with someone who has crabs Sharing personal items (towels, bedding) that are infested with crabs or crab eggs	Prescription and over-the-counter medication (creams, lotions, or shampoos) Must thoroughly clean bedding, towels, etc.
Genital Warts (HPV)	3-6 months after infection Sometimes can't be seen at all Small, bumpy growth(s) on or around sex organs or anus Can be inside the vagina Can vary in color and size Can cause cervical cancer in women. Annual PAP test recommended to detect warts	Skin-to-skin sexual contact (vaginal, oral, anal) with a person who has genital warts Mother can spread to infant during childbirth Much still unknown about the virus 70-85% sexually active people are infected	A doctor can remove genital warts, but they can grow back
Gonorrhea	2-10 days after infection Sometimes NO SYMPTOMS Pain or burning with urination Discharge from penis or vagina Severe sore throat Males are more likely to have symptoms than females Can lead to Pelvic Inflammatory Disease (PID) if left untreated	Sexual contact (vaginal, anal, oral) Mother can pass it to an infant during childbirth	Antibiotics prescribed by a doctor (pills or shot) Your sex partner(s) must also be treated
HIV/AIDS	It could take 10-12 years to get symptoms after infection Symptoms include fever, night sweats, unexplained weight loss, diarrhea, swollen lymph glands	Sexual contact (vaginal, anal, oral) Exchange of blood, semen, pre-cum, and vaginal fluids with an infected person Sharing needles Mother can pass it to an infant before or during birth or through breast milk	There is NO CURE Prescription medications help to slow down HIV & treat infections caused by a weakened immune system
Non-Gonococcal Urethritis (NGU)	2-21 days after infection Discharge from penis or vagina Pain or burning during urination Lower abdominal pain Sometimes NO SYMPTOMS	Sexual contact (vaginal, anal, oral) Mother can pass it to an infant during childbirth.	Cured with antibiotic medications prescribed by a doctor. Your sex partner(s) must also be treated.
Syphilis	1 st stage: 1-12 wks after infection, Painless sore anywhere in genital area or mouth 2 nd stage: 2-12 wks after sore has healed, skin rashes and flu-like symptoms may appear 3 rd stage: latency (no symptoms)	Sexual contact (vaginal, anal, oral) with someone who has syphilis. Mother can pass it to an infant during childbirth	Prescription antibiotic medication Sexual partners must be tested and treated if infected
Vaginitis (BV, Yeast, Trichomoniasis)	Usually itching, pain, burning in vagina Unusual discharge from vagina that might have an odor	Can be spread during sexual contact Men can carry germs without having symptoms	Yeast: Prescription or over-the-counter medication Trich: Prescription medication. Partners need to be treated BV: Prescribed antibiotics